























Medication	Photo	Tablets Or Capsules Per day	Possible Side Effects
B6 (Pyridoxine) 25mg No imprint	 25mg		Feeling of slowness (sluggishness). Lack of feeling in the hands and feet. Stomach ache.
B6 (Pyridoxine) 50mg No imprint	 50mg		Feeling of slowness (sluggishness). Lack of feeling in the hands and feet. Stomach ache.
Ethambutol 100mg white tablet "VP 11"	 100mg		Call your doctor if you have any of these side effects: Any vision changes such as blurring, color blindness, trouble seeing. Eye pain. Swelling of face, lips, or throat. Large bumps on skin. Wheezing or trouble breathing. Lack of feeling, pain, or tingling in hands or feet. Joint pain. Fever or chills. Bumps on skin that is red or itchy. Feel like throwing up. Stomach pain. Headache. Dizziness. Do not want to eat.
Ethambutol 400mg white tablet "VP 14"	 400mg		Call your doctor if you have any of these side effects: Any vision changes such as blurring, color blindness, trouble seeing. Eye pain. Swelling of face, lips, or throat. Large bumps on skin. Wheezing or trouble breathing. Lack of feeling, pain, or tingling in hands or feet. Joint pain. Fever or chills. Bumps on skin that is red or itchy. Feel like throwing up. Stomach pain. Headache. Dizziness. Do not want to eat.
Isoniazid 100mg white tablet "066/100"	 100mg		Call your doctor if you have any of these side effects: Yellow skin or eyes. Dark, coffee or tea-colored urine. Very bad stomach pain. Feel like throwing up. Weakness. Hard time seeing or eye pain. Fever. Diarrhea. Bumps on skin that is red or itchy. Avoid alcohol with this medication.
Isoniazid 300mg white tablet "Westward 261"	 300mg		Call your doctor if you have any of these side effects: Yellow skin or eyes. Dark, coffee or tea-colored urine. Very bad stomach pain. Feel like throwing up. Weakness. Hard time seeing or eye pain. Fever. Diarrhea. Bumps on skin that is red or itchy. Avoid alcohol with this medication.
Pyrazinamide 500mg white tablet "VP 012"	 500mg		Call your doctor if you have any of these side effects: Bumps on skin is very red, very itchy, or very large. Pain or swelling in the joints. Yellowing of the skin or eyes. Fever or chills. Dark urine. Feeling like throwing up. Unusual tiredness or do not want to eat.
Rifadin (Rifampin) 150mg red capsule "Rifadin 150"	 150mg		Call your doctor if you have any of these side effects: Feeling like throwing up. Yellowing of skin or eyes. Fever. Lack of feeling, pain, or tingling in arms or legs. Feeling weak or having a hard time standing straight. Hard time seeing things or eye pain. Bloody urine. Coffee or tea colored urine. Unusual bleeding or bruising. Bumps on skin that is red or itchy. Best if taken on an empty stomach.
Rifadin (Rifampin) 300mg red capsule "Rifadin 300"	 300mg		Call your doctor if you have any of these side effects: Feeling like throwing up. Yellowing of skin or eyes. Fever. Lack of feeling, pain, or tingling in arms or legs. Feeling weak or having a hard time standing straight. Hard time seeing things or eye pain. Bloody urine. Coffee or tea colored urine. Unusual bleeding or bruising. Bumps on skin that is red or itchy. Best if taken on an empty stomach.
Rifamate™ red capsule "Rifamate"	 (rifampin 300mg, isoniazid 150mg)		This medication has Isoniazid and Rifadin (Rifampin). Please read the side effects for each one.
Rifater™ pink coated tablet "Rifater"	 (rifampin 120mg, isoniazid 50mg, pyrazinamide 300mg)		This medication has Isoniazid, Rifadin (Rifampin), and Pyrazinamide. Please read the side effects for each one.

Health Center

PHN: _____ M.D.: _____

County of Los Angeles Public Health. Tuberculosis Control Program.

Medicina	Fotografia	Tabletas o Capsulas al dia	Efectos Secundarios Posible
B6 (Pyridoxine) 25mg No imprint	 25mg		Torpeza. Adormecimiento de pies y manos. Malestar estomacal.
B6 (Pyridoxine) 50mg No imprint	 50mg		Torpeza. Adormecimiento de pies y manos. Malestar estomacal.
Ethambutol 100mg white tablet "VP 11"	 100mg		Consulte con el doctor si observa cualquiera de estos efectos secundarios: Cualquier cambio de la vista cómo visión borrosa, ceguera de colores, problemas para ver. Dolor de ojos. Hinchazon en la cara, labios, garganta. Ronchas. Silbidos o problemas de respiración. Adormecimiento, dolor, o cosquilleo en pies o manos. Dolor en las coyunturos. Fiebre escalofrios. Salpullido en la piel. Dolor en la cabeza. Falta de apetito.
Ethambutol 400mg white tablet "VP 14"	 400mg		Consulte con el doctor si observa cualquiera de estos efectos secundarios: Cualquier cambio de la vista cómo visión borrosa, ceguera de colores, problemas para ver. Dolor de ojos. Hinchazon en la cara, labios, garganta. Ronchas. Silbidos o problemas de respiración. Adormecimiento, dolor, o cosquilleo en pies o manos. Dolor en las coyunturos. Fiebre escalofrios. Salpullido en la piel. Dolor en la cabeza. Falta de apetito.
Isoniazid 100mg white tablet "066/100"	 100mg		Consulte con el doctor si observa cualquiera de estos efectos secundarios: ojos o piel amarilla. Orina oscura o de color café. Dolor estomacal severo, nausea y vomitó. Debilidad. Visión borrosa, dolor de ojos. Fiebre. Salpullido o diarrea. No consuma bebidas alcoholicas mientras este tomando esta medicina.
Isoniazid 300mg white tablet "Westward 261"	 300mg		Consulte con el doctor si observa cualquiera de estos efectos secundarios: ojos o piel amarilla. Orina oscura o de color café. Dolor estomacal severo, nausea y vomitó. Debilidad. Visión borrosa, dolor de ojos. Fiebre. Salpullido o diarrea. No consuma bebidas alcoholicas mientras este tomando esta medicina.
Pyrazinamide 500mg white tablet "VP 012"	 500mg		Consulte con el doctor si observa cualquiera de estos efectos secundarios: Salpullido, picazón severa o ronchas. Dolor o hinchazon en las coyunturas. Ojos y piel amarilla. Fiebre o escalofrios. Orina oscura. Nausea vomitó cansansio. Falta de apetito.
Rifadin (Rifampin) 150mg red capsule "Rifadin 150"	 150mg		Consulte con el doctor si observa cualquiera de estos efectos secundarios: Nausea y vomitó. Ojos y piel amarilla. Fiebre. Adormecimiento, dolor, o cosquilleo en brazos o piernas. Debilidad inestable. Visión borrosa o dolor de ojos. Sangre en la orina o color café. Sangrado inusual. Salpullido o comezón. Tome con estómago vacío.
Rifadin (Rifampin) 300mg red capsule "Rifadin 300"	 300mg		Consulte con el doctor si observa cualquiera de estos efectos secundarios: Nausea y vomitó. Ojos y piel amarilla. Fiebre. Adormecimiento, dolor, o cosquilleo en brazos o piernas. Debilidad inestable. Visión borrosa o dolor de ojos. Sangre en la orina o color café. Sangrado inusual. Salpullido o comezón. Tome con estómago vacío.
Rifamate™ red capsule "Rifamate"	 (rifampin 300mg, isoniazid 150mg)		Esta medicina es combinación de Isoniazid y Rifampin. Consulte por favor la información individual de cada ingrediente.
Rifater™ pink coated tablet "Rifater"	 (rifampin 120mg, isoniazid 50mg, pyrazinamide 300mg)		Esta medicina es combinación de Isoniazid, Rifampin y Pyrazinamide. Consulte por favor la información individual de cada ingrediente.

Health Center

PHN: _____

M.D.: _____